

# Chlorophyll Complex Ointment™

2340

Please Copy for Your Patients

## Chlorophyll Complex Ointment Contains Chlorophyll and *Tillandsia usneoides* to Help Maintain Healthy Skin

The ingredients contained in Chlorophyll Complex Ointment have a long history of use to keep your skin soft and healthy.†

### How Chlorophyll Complex Ointment Keeps You Healthy

#### Maintains healthy skin

*Tillandsia usneoides*, an ingredient of Chlorophyll Complex Ointment, is reported to be used extensively in South America. Chlorophyll Complex Ointment contains naturally-occurring vitamin C from *Tillandsia usneoides*. Vitamin C is well documented for its important role in the formation of collagen, necessary for skin health.

Some evidence suggests that the observed medical effects of chlorophyll in plants are the result of interaction between different components of the whole plant. Other phytochemicals present in Chlorophyll Complex Ointment that may work in combination with the chlorophylls include carotenoids, flavonoids, indole, isothiocyanate, polyphenolic compounds, and protease inhibitors.†



**Introduced in:**

1949

**Content:**

Net Weight 18 grams

Chlorophyll Complex Ointment™ 2340



800-558-8740 • [www.standardprocess.com](http://www.standardprocess.com)

† These statements have not been evaluated by the Food & Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

# Chlorophyll Complex Ointment™

## What Makes Chlorophyll Complex Ointment Unique Unique Product Attributes

### Ingredients are derived from whole-food sources

- Contains crude chlorophyll extract from alfalfa, pea vine, buckwheat, and *Tillandsia usneoides*
- Provides vitamin K, lutein, and other various vitamins and minerals

### Certified Organic Farming

A healthy ecosystem is created by using organic farming techniques, such as rotating crops, fertilizing the soil with nutrient-rich cover crops and by-products from our processing, practicing strict weed control standards, and continually monitoring the health of our plants

- Assures the soil is laden with minerals and nutrients
- Ensures plants are nutritionally complete and free from synthetic pesticides

### Unique Processing

Upon harvesting, nutrient-rich plants are immediately washed and promptly processed

- Preserves nutritional integrity

### Exclusive low-temperature, high-vacuum drying technique

- Preserves the enzymatic vitality and nutritional potential of ingredients

### Not disassociated into isolated components

- The nutrients in Chlorophyll Complex Ointment are processed to remain intact, complete nutritional compounds

Degreed microbiologists and chemists in our on-site laboratories constantly conduct bacterial and analytical tests on raw materials, product batches, and finished products

- Ensures consistent quality and safety

### Vitamin and mineral analyses validate product content and specifications

- Assures high-quality essential nutrients are delivered

### Whole Food Philosophy

Dr. Lee challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature—in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists—known and unknown—bioactivity is markedly enhanced over synthetic nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to a synthetic or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.

**Ingredients:** Soybean lecithin, crude chlorophyll extract (from alfalfa, pea (vine), buckwheat, and *Tillandsia usneoides*), lard, lanolin, beeswax, flaxseed oil, and bovine orchid glandular extract.

**Suggested Use:** To be applied locally as an emollient.

**Special Information:** For external use only.  
Sold to health care professionals.

Studies on nutrients generally use large doses and these studies, some of which are cited below, are the basis for much of the information we provide you in this publication about whole food ingredients.

Arny N.P. "Spanish Moss and Ball Moss." University of Florida. Cooperative Extension Service Institute of Food and Agricultural Sciences. Online. 17 May 2000.

Chernomorsky S., et al. 1999. Effect of dietary chlorophyll derivatives on mutagenesis and tumor cell growth. *Teratog Carcinog Mutagen* 19(5): 313-322.

Costa M., Di Stasi L.C., Kirizawa M., et al. 1989. Screening in mice of some medicinal plants used for analgesic purposes in the state of Sao Paulo. Part II. *J Ethnopharmacol* 27(1-2): 25-33.

Di Padova C., Bosisio E., Cighetti G., et al. 1982. 3-Hydroxy-3-methylglutaric acid (HMGA) reduces dietary cholesterol induction of saturated bile in hamster. *Life Sci* 30(22): 1907-1914.

di Stasi L.C., Costa M., Mendacoli S.L., et al. 1988. Screening in mice of some medicinal plants used for analgesic purposes in the state of Sao Paulo. *J Ethnopharmacol* 24(2-3): 205-211.

Duke J.A. 1992. *Handbook of phytochemical constituents of GRAS herbs and other economic plants*. Boca Raton, FL: CRC Press, Inc: 28.

Duke J. 2000. *Phytochemical and Ethnobotanical Database, USDA – ARS – NGRIL*. Beltsville Agricultural Research Center, Beltsville, Maryland. Online. 22 May 2000.

Francesconi M., Galzigna L., Plebani M., et al. 1987. Effect of 3-hydroxy-3-methylglutarate treatment on plasma ketone bodies, triglycerides, and HDL-cholesterol in streptozotocin-diabetic rats. *Clin Biochem* 20(3): 201-205.

Furukawa E., et al. 1998. A 13-week subchronic oral toxicity study of chlorophyll in F344 rats. *Kokuritsu Iyakuin Shokuhin Eisei Kenkyusho Hokoku* (116): 107-112.

Gentile J.M., et al. 1991. The metabolic activation of 4-nitro-o-phenylenediamine by chlorophyll-containing plant extracts: the relationship between mutagenicity and antimutagenicity. *Mutat Res* 250(1-2): 79-86.

Gregory R. 1989. *Biochemistry of Photosynthesis*. Chichester, England: John Wiley and Sons, Ltd: 10.

Hajri A., et al. 1999. Human pancreatic carcinoma cells are sensitive to photodynamic therapy in vitro and in vivo. *Br J Surg* 86(7): 899-906.

Kaufman P.B., et al. 1999. *Natural products from plants*. Boca Raton, FL: CRC Press: 61.

Labate M.E., Dam R. 1980. Effect of 3-hydroxy-3-methylglutaric acid on cholesterol metabolism in female Japanese quail. *Poult Sci* 59(2): 383-389.

©2005 Standard Process Inc. All rights reserved. 12/05

Lea P.J., Leegood R.C., eds. 1999. *Plant biochemistry and molecular biology*. West Sussex, England: John Wiley and Sons, Ltd: 233.

Lupien P.J., Moorjani S., Brun D., et al. 1979. Effects of 3-hydroxy-3-methylglutaric acid on plasma and low-density lipoprotein cholesterol levels in familial hypercholesterolemia. *J Clin Pharmacol* 19(2-3): 120-126.

Moorjani S., Lupien P.J. 1977. Effect in vitro of 3-hydroxy-3-methylglutaric acid on the synthesis of mevalonate and its precursors. *Arch Int Physiol Biochim* 85(1): 1-10.

Negishi T., et al. 1989. Inhibitory effect of chlorophyll on the genotoxicity of 3-amino-1-methyl-5H-pyrido[4,3-b]indole (Trp-P-2). *Carcinogenesis* 10(1): 145-149.

Negishi T., et al. 1997. Antigenotoxic activity of natural chlorophylls. *Mutat Res* 376(1-2): 97-100.

Sarkar D., et al. 1996. Chlorophyll and chromosome breakage. *Mutat Res* 360(3): 187-191.

Savoie L.L., Lupien P.J. 1975. Organ distribution of 3-hydroxy-3-methylglutaric acid, a potential anticholesterolemic agent. *Can J Physiol Pharmacol* 53(4): 638-643.

Savoie L.L., Lupien P.J. 1975. Preliminary toxicological investigations of 3-hydroxy-3-methylglutaric acid (HMGA). I: Study of acute toxicity and of teratogenic activity in rats and mice. *Arzneimittelforschung* 25(8): 1284-1286.

Van Niekerk J.L., Hendriks T., Gevers J.A., et al. 1984. The lipid-lowering effects of 3-hydroxy-3-methylglutaric acid and bile acid drainage in WHHL rabbits. *Clin Sci* 67(4): 439-444.

Witherup K.M., McLaughlin J.L., Judd R.L., et al. 1995. Identification of 3-hydroxy-3-methylglutaric acid (HMGA) as a hypoglycemic principle of Spanish moss (*Tillandsia usneoides*). *J Nat Prod* 58(8): 1285-1290.

Yousufzai S.Y., Siddiqi M. 1976. 3-Hydroxy-3-methylglutaric acid and experimental atherosclerosis in rats. *Experientia* 32(8): 1033-1034.

Yousufzai S.Y., Siddiqi M. 1976. 3-Hydroxy-3-methylglutaric acid and triton-induced hyperlipidemia in rats. *Experientia* 32(9): 1178-1179.

Yousufzai S.Y., Siddiqi M. 1977. Serum and liver lipid responses to 3-hydroxy-3-methylglutaric acid in rats on different carbohydrate diets. *Lipids* 12(3): 262-266.

Yousufzai S.Y., Siddiqi M. 1977. Tissue lipid responses to 3-hydroxy-3-methylglutaric acid with different dietary fats. *Lipids* 12(3): 258-261.

Yousufzai S.Y., Siddiqi M., Abdullah A.K. 1976. Protective effects of 3-hydroxy-3-methylglutaric acid in alcohol-induced lipemia. *Lipids* 11(7): 526-529.